

File: References/word count:10,258/page count:44

;a References

;b Introduction

Bodnar L, Wisner K: Nutrition and depression. Biological Psychiatry 2005;58:679-685.

Bogner H, Morales K, Post E, et al: Diabetes, depression, and death. Diabetes Care 2007;30:3005-3010.

Bonnet F, Irving K, Terra J, et al: Depressive symptoms are associated with unhealthy lifestyles in hypertensive patients with the metabolic syndrome. Journal of Hypertension 2005;23:611-617.

Buscemi D, Kumar A, Nugent R, et al: Short sleep times predict obesity in internal medicine clinic patients. Journal of Clinical Sleep Medicine 2007;3:681-688.

Christensen K, McGue M, Petersen I, et al: Exceptional longevity does not result in excessive levels of disability. Proceedings of the National Academy of Sciences 2008;105:13274-13279.

Christensen L, Redig C: Effect of meal composition on mood. Behavioral Neurology 1993;107:346-353.

Christensen L, Somers S: Adequacy of the dietary intake of depressed individuals. Journal of the American College of Nutrition 1994;13:597-600.

Depression linked to bone-thinning in premenopausal women. Science Daily 2007;November 27th.

Gibson E: Emotional influences on food choice. Physiology & Behavior 2006;89:53-61.

Kizilay P: Predictors of depression in women. Women's Health 1992;27:983-993.

Klem M, Wing R, Lang W, et al: Does weight loss maintenance become easier over time? Obesity Research 2000;8:438-444.

Klem M, Wing R, McGuire M, et al: Psychological symptoms in individuals successful at long-term maintenance of weight loss. Health Psychology 1998;17:336-345.

Klem M, Wing R, McGuire M, et al: A descriptive study of individuals successful at long-term maintenance of substantial weight loss. American Journal of Clinical Nutrition 1997;66:239-246.

Liang S, Byers D, Irwin L: Chronic mild stressors and diet affect gene expression differently in male and female rats. Journal of Molecular Neuroscience 2007;33:189-200.

Liu C, Xie B, Chou C, et al: Perceived stress, depression and food consumption frequency in the college students of China Seven Cities. Physiology & Behavior 2007;92:748-754.

Lombard C: What is the role of food in preventing depression and improving mood, performance and cognitive function? Medical Journal of Australia 2000;173:S104-S105.

McGuire M, Wing R, Klem M, et al: What predicts weight regain in a group of successful weight losers? Journal of Consultation and Clinical Psychology 1999;67:177-185.

Petry N, Barry D, Pietrzak R, et al: Overweight and obesity are associated with psychiatric disorders. Psychosomatic Medicine 2008;70:288-297.

Rogers P: A healthy body, a healthy mind: Long-term impact of diet on mood and cognitive function. Proceedings of the Nutrition Society 2001;60:135-143.

Severus W, Littman A, Stoll A: Omega-3 fatty acids, homocysteine, and the increased risk for cardiovascular mortality in major depressive disorder. Harvard Review of Psychiatry 2001;9:280-293.

Simon G, Ludman E, Linde J, et al: Association between obesity and depression in middle-age women. General Hospital Psychiatry 2008;1:32-39.

Steptoe A, O'Donnell K, Badrick E, et al: Neuroendocrine and inflammatory factors associated with positive affect in healthy men and women. American Journal of Epidemiology

2008;167:96-102.

Wadden T, Steen S, Wingate B, et al: Psychosocial consequences of weight reduction. American Journal of Clinical Nutrition 1996;63:S461-S465.

Whitmer R, Gustafson D, Barrett-Connor E, et al: Central obesity and increased risk of dementia more than three decades later. Neurology 2008;71:1057-1064.

Williams A, Cotter A, Sabina A, et al: The role of vitamin B6 as treatment for depression. Family Practice 2005;22:532-537.

Wurtman J: Depression and weight gain: The serotonin connection. Journal of Affective Disorders 1993;29:183-192.

Wurtman R, Wurtman J: Carbohydrates and depression. Scientific American. January 1989.

Yoshioka M, Ito K, Harada R, et al: Association among nutritional status, mood and body composition. FASEB Journal 2000;14:A254.

Ziegelstein R, Fauerbach J, Stevens S, et al: Patients with depression are less likely to follow recommendations to reduce cardiac risk during recovery from a myocardial infarction. Archives of Internal Medicine 2000;160:1818-1823.

;b Chapter 1

Barberger O, Gateau P, Raffaitin C, et al: Dietary patterns and risk of dementia. Neurology 2007;69:1921-1930.

Crawley J: The role of galanin in feeding behavior. Neuropeptides 1999;33:369-375.

Jahng J, Kim J, Kim H, et al: Chronic food restriction in young rats results in depression and anxiety-like behaviors with decreased expression of serotonin reuptake transporter. Brain Research 2007;1150:100-107.

Kruger J, Blanck H, Gillespie C: Dietary practices, dining out behavior, and physical activity correlates with weight loss maintenance. Preventing Chronic Disease 2008;5:A11.

Lawrence C, Baudoin F, Luckman S: Centrally administered galanin-like peptide modifies food intake in the rat: A comparison with galanin. Journal of Neuroendocrinology 2002;14:853-860.

Leibowitz S: Differential functions of hypothalamic galanin cell groups in the regulation of eating and body weight. Annals of the New York Academy of Sciences 1998;863:206-220.

Nothlings U, Schulze M, Welkert C, et al: Intake of vegetables, legumes, and fruit, and risk for all-cause, cardiovascular and cancer mortality in a European diabetic population. Journal of Nutrition 2008;138:775-781.

Parrott M, Greenwood C: Dietary influences on cognitive function with aging: From high-fat diets to healthful eating. Annals of the New York Academy of Sciences 2007;1114:389-397.

Raynor H, Jeffery R, Phelan S, et al: Amount of food group variety consumed in the diet and long-term weight loss maintenance. Obesity Research 2005;13:883-890.

Sanchez-Villegas A, Henriquez P, Bes-Rastrollo, et al: Mediterranean diet and depression. Public Health Nutrition 2006;9:1104-1109.

Small G, Silverman D, Siddarth P, et al: Effects of a 14-day healthy longevity lifestyle program on cognition and brain function. American Journal of Geriatric Psychiatry 2006;14:538-545.

Smith A, Clark R, Gallagher J: Breakfast cereal and caffeinated coffee: Effects on working memory, attention, mood, and cardiovascular function. Physiology & Behavior 1999;67:9-17.

Spring B, Chiodo J, Harden M, et al: Psychobiological effects of carbohydrates. Journal of Clinical Psychiatry 1989;50:27-33.

Tucker K, Hallfrisch J, Qiao N, et al: The combination of high fruit and vegetable and low saturated fat intakes is more protective against mortality in aging men than is either alone.

Journal of Nutrition 2005;135:556-561.

Wang J, Akabayashi A, Yu H, et al: Hypothalamic galanin: Control by signals of fat metabolism.

Brain Research 1998;804:7-20.

Wilding J: Neuropeptides and appetite control. Diabetes Medicine 2002;19:619-627.

;b Chapter 2

Benton D, Maconie A, Williams C: The influence of the glycaemic load of breakfast on the behaviour of children in school. Physiology & Behavior 2007;92:717-724.

Bertrais S, Luque M, Preziosi P, et al: Contribution of ready-to-eat cereals to nutrition intakes in French adults and relations with corpulence. Annals of Nutrition & Metabolism 2000; 44:249-255.

Cho S, Dietrich M, Brown C, et al: The effect of breakfast type on total daily energy intake and body mass index: results from the third National Health and Nutrition Examination Survey (NHANES III). Journal of the American College of Nutrition 2003;22:296-302.

Ciok J, Dolna A: Carbohydrates and mental performance: The role of glycemic index of food products. Pol Merkur Lekarski 2006;20:367-370.

Elfhag K, Rossner S: Who succeeds in maintaining weight loss? A conceptual review of factors associated with weight loss maintenance and weight regain. Obesity Review 2005;6:67-85.

Farshchi H, Taylor M, Macdonald I: Deleterious effects of omitting breakfast on insulin sensitivity and fasting lipid profiles in healthy lean women. American Journal of Clinical Nutrition 2005;81:388-396.

Good, C, Holschuh N, Albertson A, et al: The relationship between whole grain consumption and body mass index in adult men and women: data from the USDA's Continuing Survey of

Food Intakes by Individuals, Pyramid Servings Data, 1994-96. Abstract; presented at FASEB, April 2004.

Greenwood C, Kaplan R, Hebblethwaite S, et al: Carbohydrate-induced memory impairment in adults with Type 2 diabetes. Diabetes Care 2003;26:1961-1966.

Holt S, Delargy H, Lawton C, et al: The effects of high-carbohydrate vs high-fat breakfasts on feelings of fullness and alertness, and subsequent food intake. International Journal of Food Science and Nutrition 1999;50:15-28.

Ingwersen J, Defeyter M, Kennedy D, et al: A low glycaemic index breakfast cereal preferentially prevents children's cognitive performance from declining throughout the morning. Appetite 2007;49:240-244.

Keim N, Van Loan M, Horn W, et al: Weight loss is greater with consumption of large morning meals and fat-free mass is preserved with large evening meals in women on a controlled weight reduction regimen. Journal of Nutrition 1997;127:75-82.

Levine A, Tallman J, Grace M, et al: Effect of breakfast on short-term food intake. American Journal of Clinical Nutrition 1989;50:1303-1307.

Liu S, Willett W, Manson J, et al: Relation between changes in intakes of dietary fiber and grain products and changes in weight and development of obesity among middle-aged women. American Journal of Clinical Nutrition 2003; 78:920-927.

Ma Y, Bertone E, Stanek E, et al: Association between eating patterns and obesity in a free-living U.S. adult population. American Journal of Epidemiology 2003; 158:85-92.

Mahoney C, Taylor H, Kanarek R, et al: Effect of breakfast composition on cognitive processes in elementary school children. Physiology & Behavior 2005;85:635-645.

Nettleton J, Steffen L, Ni H, et al: Dietary patterns and risk of incident type 2 diabetes in the

Multi-Ethnic Study of Atherosclerosis (MESA). Diabetes Care 2008;31:1777-1782.

Newby P, Mueller D, Hallfrisch J, et al: Dietary patterns and changes in body mass index and waist circumference in adults. American Journal of Clinical Nutrition 2003;77:1417-1425.

Nilsson A, Ostman E, Granfeldt Y, et al: Effect of cereal test breakfasts differing in glycemic index and content of indigestible carbohydrates on daylong glucose tolerance in healthy subjects. American Journal of Clinical Nutrition 2008;87:645-654.

Pasman W, Blokdijk V, Bertina F, et al: Effect of two breakfasts, different in carbohydrate composition, on hunger and satiety and mood in healthy men. International Journal of Obesity 2003;27:663-668.

Schlundt D, Hill J, Sbrocco T, et al: The role of breakfast in the treatment of obesity: a randomized clinical trial. American Journal of Clinical Nutrition 1992;55:645-651.

Smith A: Stress, breakfast cereal consumption and cortisol. Nutritional Neuroscience 2002;5:141-144.

Smith A, Clark R, Gallagher J: Breakfast cereal and caffeinated coffee: Effects on working memory, attention, mood, and cardiovascular function. Physiology & Behavior 1999;67:9-17.

Timlin M, Pereira M, Story M, et al: Breakfast eating and weight change in a 5-year prospective analysis of adolescents. Pediatrics 2008;121:e638-e645.

Williams P. Breakfast and the diets of Australian adults: An analysis of data from the 1995 National Nutrition Survey. International Journal of Food Sciences and Nutrition 2005; 56(1):65-79.

Wyatt H, Grunwald G, Mosca C, et al: Long-term weight loss and breakfast in subjects in the National Weight Control Registry. Obesity Research 2002;10:78-82.

;b Chapter 3

Afaghi A, O'Connor H, Chow C: High-glycemic index carbohydrate meals shorten sleep onset.

American Journal of Clinical Nutrition 2007;85:426-430.

Andlauer W, Furst P: Antioxidative power of phytochemicals with special reference to cereals.

Cereal Foods World 1998;43:356-360.

Bendich A: The potential for dietary supplements to reduce premenstrual syndrome (PMS)

symptoms. Journal of the American College of Nutrition 2000;19:3-12.

Benton D: Carbohydrate ingestion, blood glucose, and mood. Neuroscience and Biobehavioral

Reviews 2002;26:293-308.

Christensen L, Redig C: Effect of meal composition on mood. Behavioral Neuroscience

1993;107:346-353.

Cleveland L, Moshfegh A, Albertson A, et al: Dietary intake of whole grains. Journal of the

American College of Nutrition 2000;19:S331-S338.

Cross G, Marley J, Miles H, et al: Changes in nutrient intake during the menstrual cycle of

overweight women with premenstrual syndrome. British Journal of Nutrition 2001;85:475-482.

Docherty J, Sack D, Roffman M, et al: A double-blind, placebo-controlled, exploratory trial of

chromium picolinate in atypical depression: Effect on carbohydrate cravings. Journal of

Psychiatric Practice 2005;11:302-314.

Dye L, Warner P, Bancroft J: Food craving during the menstrual cycle and its relationship to

stress, happiness of relationship and depression: A preliminary enquiry. Journal of Affective

Disorders 1995;34:157-164.

Foster-Powell K, Holt S, Brand-Miller J: International table of glycemic index and glycemic load

values. American Journal of Clinical Nutrition 2002;76:5-56.

Jacobs D, Meyer K, Kushi L, et al: Whole-grain intake may reduce the risk of ischemic heart disease death in postmenopausal women. American Journal of Clinical Nutrition 1998;68:248-257.

Halyburton A, Brinkworth G, Wilson C, et al: Low- and high-carbohydrate weight-loss diets have similar effects on mood, but not cognitive performance. American Journal of Clinical Nutrition 2007;86:580-587.

Klaassen T, Riedel W, van Someren A, et al: Mood effects of 24 hour tryptophan depletion in healthy, first-degree relatives of patients with affective disorders. Biology Psychiatry 1999;46:480-497.

Katcher H, Legro R, Kunselman A, et al: The effects of a whole grain-enriched Hypocaloric diet on cardiovascular disease risk factors in men and women with metabolic syndrome. American Journal of Clinical Nutrition 2008;87:79-90.

Liu S: Intake of refined carbohydrates and whole grain foods in relation to risk of type 2 diabetes mellitus and coronary heart disease. Journal of the American College of Nutrition 2002;21:298-306.

Ludwig D: Dietary glycemic index and the regulation of body weight. Lipids 2003;38:117-121.

Ludwig D: Dietary glycemic index and obesity. Journal of Nutrition 2000;130:S280-S283.

Markus C: Effects of carbohydrates on brain tryptophan availability and stress performance. Biological Psychology 2007;76:83-90.

McKeown N, Meigs J, Liu S, et al: Whole-grain intake is favorably associated with metabolic risk factors for type 2 diabetes and cardiovascular disease in the Framingham Offspring Study. American Journal of Clinical Nutrition 2002;76:390-398.

Murakami K, Sasaki S, Takahashi Y, et al: Dietary glycemic index is associated with decreased

premenstrual symptoms in young Japanese women. Nutrition 2008;March 21st.

Muthayya S, Thomas T, Srinivasan K, et al: Consumption of a mid-morning snack improves memory but not attention in school children. Physiology & Behavior 2007;90:142-150.

Newby P, Maras J, Bakun P, et al: Intake of whole grains, refined grains, and cereal fiber measured with 7-d diet records and associations with risk factors for chronic disease. American Journal of Clinical Nutrition 2007;86:1745-1753.

Papanikolaou Y, Palmer H, Binns M, et al: Better cognitive performance following a low-glycemic compared with a high-glycemic index carbohydrate meal in adults with type 2 diabetes. Diabetologia 2006;49:855-862.

Pins J, Geleva D, Keenan J, et al: Do whole-grain oat cereals reduce the need for antihypertensive medications and improve blood pressure control? Journal of Family Medicine 2002;51:353-359.

Reed S, Levin F, Evans S: Changes in mood, cognitive performance and appetite in the late luteal and follicular phases of the menstrual cycle in women with and without PMDD (premenstrual dysphoric disorder). Hormones and Behavior 2008;54:185-193.

Slavin J: Mechanisms for the impact of whole grain foods on cancer risk. Journal of the American College of Nutrition 2000;19:S300-S307.

Spring B, Chiodo J, Harden M, et al: Psychobiological effects of carbohydrates. Journal of Clinical Psychiatry 1989;50 (suppl):27-33.

Spring B, Schneider K, Smith M, et al: Abuse potential of carbohydrates for overweight carbohydrate cravers. Psychopharmacology 2008;197:637-647.

Sweat V, Starr V, Bruehl H, et al: C-reactive protein is linked to lower cognitive performance in overweight and obese women. Inflammation 2008;31:198-207.

Tanke M, Alserda E, Doornbos B, et al: Low tryptophan diet increases stress-sensitivity, but does not affect habituation in rats. Neurochemistry International 2008;52:272-281.

Truswell A: Cereal grains and coronary heart disease. European Journal of Clinical Nutrition 2002;56:1-14.

Uchida S, Umeeda H, Kitamoto A, et al: Chronic reduction in dietary tryptophan leads to selective impairment of contextual fear memory in mice. Brain Research 2007;1149:149-156.

van der Stelt H, Broersen L, Olivier B, et al: Effects of dietary tryptophan variations on extracellular serotonin in the dorsal hippocampus of rats. Psychopharmacology 2004;172:137-144.

Wurtman J: Depression and weight gain: The serotonin connection. Journal of Affective Disorders 1993;29:183-192.

;b Chapter 4

www.ers.usda.gov/AmberWaves/April03/Indicators/behinddata.htm

Ackroff K, Sclafani A: Sucrose-induced hyperphagia and obesity in rats fed a macronutrient self-selection diet. Physiology & Behavior 1988;44:181-187.

Adam T, Epel E: Stress, eating and the reward system. Physiology & Behavior 2007;91:449-458.

Avena N, Bocarsly M, Rada P, et al: After daily binging on a sucrose solution, food deprivation induces anxiety and accumbens dopamine/acetylcholine imbalance. Physiology & Behavior 2008;94:309-315.

Avena N, Long K, Hoebel B: Sugar-dependent rats show enhanced responding for sugar after abstinence. Physiology & Behavior 2005;84:359-362.

Barbano M, Cador M: Opioids for hedonic experience and dopamine to get ready for it.

Psychopharmacology 2007;191:497-506.

Cao D, Lu H, Lewis T, et al: Intake of sucrose-sweetened water induces insulin resistance and exacerbates memory deficits and amyloidosis in a transgenic mouse model of Alzheimer disease.

Journal of Biological Chemistry 2007;282:36275-36282.

Charlton K, Wolmarans P, Lombard C: Evidence of nutrient dilution with high sugar intakes in older South Africans. Journal of Human Nutrition and Dietetics 1998;11:331-343.

Chepulis L: The effect of honey compared to sucrose, mixed sugars, and a sugar-free diet on weight gain in young rats. Journal of Food Science 2007;72:S224-S229.

Chepulis L, Sarkey N: The long-term effects of feeding honey compared with sucrose and sugar-free diet on weight gain, lipid profiles, and DEXA measurements in rats. Journal of Food Science 2008;73:H1-H7.

Christensen L, Burrows R: Dietary treatment of depression. Behavior Therapy 1990;21:183-191.

Christensen L, Krietsch K, White B, et al: Impact of a dietary change on emotional distress. Journal of Abnormal Psychology 1985;94:565-579.

Dallman M, Pecoraro N, Akana S, et al: Chronic stress and obesity: A new view of “comfort food.” Proceedings of the National Academy of Sciences 2003;100:11696-11701.

Dallman M, Pecoraro N, la Fleur S: Chronic stress and comfort foods: Self medication and abdominal obesity. Brain, Behavior, and Immunity 2005;19:275-280.

Davis, J, Alexander K, Ventura E, et al: Associations of dietary sugar and glycemic index with adiposity and insulin dynamics in overweight Latino youth. American Journal Clinical Nutrition 2007;86:1331-1338.

Drewnowski A, Holden-Wiltse J: Taste responses and food preferences in obese women.

International Journal of Obesity 1992;16:639-648.

Drewnowski A, Kurth C, Holden-Wiltse J, et al: Food preferences in human obesity. Appetite 1992;18:207-221.

Emmett P, Heaton K: Is extrinsic sugar a vehicle for dietary fat? Lancet 1995;345:1537-1540.

Erianson-Albertsson C: Sugar triggers our reward system Lakartidningen 2005;102:1620-1622, 1625,1627.

Greenwood J, Standord J: Preventing or improving obesity by addressing specific eating patterns. Journal of the American Board of Family Medicine 2008;21:135-140.

Johnson R, Frary C: Choose beverages and foods to moderate your intake of sugars. Journal of Nutrition 2001;131:2766-2771.

Kranz S, Smiciklas-Wright H, Siega-Riz A, et al: Adverse effect of high added sugar consumption on dietary intake in American preschoolers. Journal of Pediatrics 2005;146:105-111.

Larsson S, Bergkvist L, Wolk A: Consumption of sugar and sugar-sweetened foods and the risk of pancreatic cancer in a prospective study. American Journal of Clinical Nutrition 2006;84:1171-1176.

Lenoir M, Serre F, Cantin L, et al: Intense sweetness surpasses cocaine reward. PloS ONE 2007;2:e698.

Lieberman H, Wurtman J, Chew B: Change in mood after carbohydrate consumption among obese individuals. American Journal of Clinical Nutrition 1986;44:772-778.

Markus R, Panhuysen G, Tuiten A, et al: Effects of food on cortisol and mood in vulnerable subjects under controllable and uncontrollable stress. Physiology & Behavior 2000;70:333-342.

Naismith D, Rhodes C: Adjustment in energy following the covert removal of sugar from the diet. Journal of Human Nutrition and Dietetics 1995;8:167-175.

Olszewski P, Levine A: Central opioids and consumption of sweet tastants. Physiology &

Behavior 2007;91:506-512.

Pecoraro N, Reyes F, Gomez F, et al: Chronic stress promotes palatable feeding, which reduces signs of stress. Endocrinology 2004;145:3754-3762.

Raben A, Vasilaras T, Moller A, et al: Sucrose compared with artificial sweeteners. American Journal of Clinical Nutrition 2002;76:721-729.

Rada P, Avena N, Hoebel B: Daily bingeing on sugar repeatedly releases dopamine in the accumbens shell. Neuroscience 2005;134:737-744.

Reid M, Hammersley R, Hill A, et al: Long-term dietary compensation for added sugar. British Journal of Nutrition 2007;97:193-203.

Salbe A, DelParigi A, Pratley R, et al: Taste preferences and body weight changes in an obesity-prone population. American Journal of Clinical Nutrition 2004;79:372-378.

Shaw G, Quach T, Nelson V, et al: Neural tube defects associated with maternal periconceptional dietary intake of simple sugars and glycemic index. American Journal of Clinical Nutrition 2003;78:972-978.

Sharma N, Okere I, Duda M, et al: High fructose diet increases mortality in hypertensive rats compared to a complex carbohydrate or high fat diet. American Journal of Hypertension 2007;20:403-409.

Swithers, S, Davidson, T: A role for sweet taste: Calorie predictive relations in energy regulation by rats. Behavioral Neuroscience 2008;122:161-173.

Spring B, Schneider K, Smith M, et al: Abuse potential for carbohydrates for overweight carbohydrate cravers. Psychopharmacology 2008;197:637-647.

Torres S, Nowson C: Relationship between stress, eating behavior, and obesity. Nutrition 2007;23:887-894.

Urias-Silvas J, Cani P, Delmee E, et al: Physiological effects of dietary fructans extracted from Agave tequilana. British Journal of Nutrition 2008;99:254-261.

Vermunt S, Pasman W, Schaafsma G, et al: Effects of sugar intake on body weight: A review. Obesity Review 2003;4:91-99.

Wansink B, Cheney M, Chan N: Exploring comfort food preferences across age and gender. Physiology & Behavior 2003;79:739-747.

Williams C, Strobino B: Childhood diet, overweight, and CVD risk factors. Preventive Cardiology 2008;11:11-20.

;b Chapter 5

Almario R, Vonghavaravat V, Wong R, et al: Effects of walnut consumption on plasma fatty acids and lipoproteins in combined hyperlipidemia. American Journal of Clinical Nutrition 2001;74:72-79.

Alper C, Mattes R: Effects of chronic peanut consumption on energy balance and hedonics. International Journal of Obesity 2002;26:1129-1137.

Andres-Lacueva C, Shukitt-Hale B, Galli R, et al: Anthocyanins in aged blueberry-fed rats are found centrally and may enhance memory. Nutrition and Neuroscience 2005;8:111-120.

Brody S, Preut R, Schommer K, et al: A randomized controlled trial of high dose ascorbic acid for reduction of blood pressure, cortisol, and subjective responses to psychological stress. Psychopharmacology 2002;159:319-324.

Burkhardt S, Tan D, Manchester L, et al: Detection and quantification of the antioxidant melatonin in Montmorency and balaton tart cherries (prunus cerasus). Journal of Agricultural and Food Chemistry 2001;49:4898-4902.

Cederroth C, Vinciguerra M, Kuhne F, et al: A phytoestrogen-rich diet increases energy expenditure and decreases adiposity in mice. Environmental Health Perspectives 2007;115:1467-1473.

Clark S: Perillyl alcohol induces c-Myc-dependent apoptosis in Bcr/Abl-transformed leukemia cells. Oncology 2006;70:13-18.

Cope M, Erdman J, Allison D: The potential role of soyfoods in weight and adiposity reduction. Obesity Research 2008;9:219-235.

Elliott A, Scheiber S, Thomas C, et al: Inhibition of glutathione reductase by flavonoids. Biochemical Pharmacology 1992;44:1603-1608.

Eagan M, Lyle R, Gunther C, et al: Effect of 1-year dairy product intervention on fat mass in young women. Obesity 2006;14:2242-2248.

Emim J, Oliveira A, Lapa A: Pharmacological evaluation of the anti-inflammatory activity of a citrus bioflavonoid, hesperidin, and the isoflavonoids, dauricin and coumestrol, in rats and mice. Journal of Pharmacy and Pharmacology 1994;46:118-122.

Engelhart M, Geerlings M, Ruitenberg A, et al: Dietary intake of antioxidants and risk of Alzheimer disease. Journal of the American Medical Association 2002;287:3223-3229.

Fernandes J, da Fonseca C, Teixeira A, et al: Perillyl alcohol induces apoptosis in human glioblastoma multiforme cells. Oncology Report 2005;13:943-947.

Flood J, Rolls B: Soup preloads in a variety of forms reduces meal energy intake. Appetite 2007;49:626-634.

Galati E, Monforte M, Kirjavainen S, et al: Biological effects of hesperidin, a citrus flavonoid: Antiinflammatory and analgesic activity. Farmaco 1994;40:709-712.

Galli R, Shukitt-Hale B, Youdim K, et al: Fruit polyphenolics and brain aging. Annals of the

New York Academy of Sciences 2002;959:128-132.

Green M, Rogers P, Elliman N, et al: Impairment of cognitive performance associated with dieting and high levels of dietary restraint. Physiology & Behavior 1994;55:447-452

Halvorsen B, Carlsen M, Phillips K, et al: Content of redox-active compounds (i.e., antioxidants) in foods consumed in the United States. American Journal of Clinical Nutrition 2006;84:95-135.

Heaney R, Davies M, Barger-Lux J: Calcium and weight: Clinical studies. Journal of the American College of Nutrition 2002;21:152S-155S.

Huang D, Ou B, Hampsch-Woodill M, et al: Development and Validation of Oxygen Radical Absorbance Capacity Assay for Lipophilic Antioxidants using Randomly Methylated Cyclodextrin as the Solubility Enhancer. Journal of Agricultural and Food Chemistry 2002;50:1815-1821.

Jaceldo-Siegl K, Sabate J, Rajaram S, et al: Long-term almond supplementation without advice on food replacement induces favourable nutrient modifications to the habitual diets of free-living individuals. British Journal of Nutrition 2004;92:533-540.

Joseph J, Denisova N, Villalobos-Molina R, et al: Oxidative stress and age-related neuronal deficits. Molecular Chemical Neuropathology 1996;28:35-40.

Joseph J, Skukitt-Hale B, Denisova N, et al: Reversals of age-related declines in neuronal signal transduction, cognitive, and motor behavioral deficits with blueberry, spinach, or strawberry dietary supplementation. Journal of Neuroscience 1999;19:8114-8121.

Joseph J, Shukitt-Hale B, Lau F: Fruit polyphenols and their effects on neuronal signaling and behavior in senescence. Annals of the New York Academy of Sciences 2007;1100:470-478.

Kim H, Park S, Han D, et al: Octacosanol supplementation increases running endurance time and improves biochemical parameters after exhaustion in trained rats. Journal of Medicinal Foods

2003;6:345-351.

Kritz-Silverstein D, Von Muhlen D, Barrett-Connor E, et al: Isoflavones and cognitive function in older women: The Soy and Postmenopausal Health in Aging (SOPHIA) study. Menopause 2003;10:189-190.

Lau F, Shukitt-Hale B, Joseph J: The beneficial effects of fruit polyphenols on brain aging. Neurobiology of Aging 2005;26 suppl:128-132.

Liao F, Shieh M, Yang S, et al: Effectiveness of a soy- based compared with a a traditional low-calorie diet on weight loss and lipid levels in overweight adults. Nutrition 2007;23:551-556.

Lin Y, Lyle R, McCabe L, et al: Dairy calcium is related to changes in body composition during a two-year exercise intervention in young women. Journal of the American College of Nutrition 2000;19:754-760.

Martin A, Cherubini A, Andres-Lacueva C, et al: Effects of fruits and vegetables on levels of vitamins E and C in the brain and their association with cognitive performance. Journal of Nutrition and Healthy Aging 2002;6:392-404.

Maskarinec G, Aylward A, Erber E, et al: Soy intake is related to a lower body mass index in adult women. European Journal of Nutrition 2008;47:138-144.

Maskarinec G, Robbins C, Riola B, et al: Three measures show high compliance in a soy intervention among premenopausal women. Journal of the American Dietetic Association 2003;103:861-866.

Mattes R: Soup and satiety. Physiology & Behavior 2005;83:739-747.

Metalla N, Braddock R, Gregory J, et al: Capillary electrophoresis and high-performance liquid chromatography determination of polyglyutamyl 5-methyltetrahydrofolate forms in citrus products. Journal of Agriculture and Food Chemistry 2005;53:2268-2274.

Morris M, Evans D, Bienias J, et al: Dietary intake of antioxidant nutrients and the risk of incident Alzheimer disease in a biracial community study. Journal of the American Medical Association 2002;287:3230-3237.

Morris M, Evans D, Tangney C, et al: Associations of vegetable and fruit consumption with age-related cognitive change. Neurology 2006;67:1370-1376.

Nagata C, Shimizu H, Takami R, et al: Serum concentrations of estradiol and dehydroepiandrosterone sulfate and soy product intake in relation to psychologic well-being in peri- and postmenopausal Japanese women. Metabolism 2000;49:1561-1564.

Orgaard A, Jensen L: The effects of soy isoflavones on obesity. Experimental Biology and Medicine 2008;June 5th.

Ou B, Hampsch-Woodill M, Prior R: Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe. Journal of Agricultural and Food Chemistry 2001;49:4619-4626.

Putnam J, Allshouse J, Kantor L: U.S. per capita food supply trends: More calories, refined carbohydrates, and fats. FoodReview 2002;Winter:2-11.

Ravaglia G, Forti P, Lucicesare A, et al: Plasma tocopherols nad risk of cognitive impairment in an elderly Italian cohort. American Journal of Clinical Nutrition 2008;87:1306-1313.

Rice-Evans C, Miller N, Bolwell P, et al: The relative antioxidant activities of plant-derived polyphenolic flavonoids. Free Radical Research 1995;22:375-383.

Rolls B, Bell E, Thorwart M: Water incorporated into a food but not served with a food decreases energy intake in lean women. American Journal of Clinical Nutrition 1999;70:448-455.

Rolls B, Fedoroff I, Guthrie J, et al: Foods with different satiating effects in humans. Appetite

1990;15:115-126.

Sabate J: Nut consumption and body weight. American Journal of Clinical Nutrition 2003;78 (3 suppl):647S-650S.

Seeram N, Bourquin L, Nair M: Degradation products of cyanidin glycosides from tart cherries and their bioactives. Journal of Agricultural and Food Chemistry 2001; 49:4924-4929.

Shukitt-Hale B, Carey A, Jenkins D, et al: Beneficial effects of fruit extracts on neuronal function and behavior in a rodent model of accelerated aging. Neurobiology of Aging 2006;July 10th.

Shukitt-Hale B, Lau F, Joseph J: Berry fruit supplementation and the aging brain. Journal of Agricultural Food Chemistry 2008;56:636-641.

Silalahi J: Anti-cancer and health protective properties of citrus fruit components. Asia Pacific Journal of Clinical Nutrition 2002;11:79-84.

Tall J, Seeram, N, Zhao C, et al: Tart cherry anthocyanins suppress inflammation-induced pain behavior in rat. Behavioural Brain Research 2004;153:181-188.

Taylor C, Hampl J, Johnston C: Low intakes of vegetables and fruits, especially citrus fruits, lead to inadequate vitamin C intakes among adults. European Journal of Clinical Nutrition 2000;54:573-578.

Tamura T, Shin Y, Buehring K, et al: The availability of folates in man: Effect of orange juice supplement on intestinal conjugase. British Journal of Haematology 1976;32:123-133.

Teegarden D, White K, Lyle R, et al: Calcium and dairy product modulation of lipid utilization and energy expenditure. Obesity 2008;April 17th.

Tsuboi H, Tatsumi A, Yamamoto K, et al: Possible connections among job stress, depressive symptoms, lipid modulation and antioxidants. Journal of Affective Disorders 2006;91:63-70.

Tsuda T, Horio F, Uchida K, et al: Dietary cyanidin 3-o-beta-D-glycoside-rich purple corn color prevents obesity and ameliorates hyperglycemia in mice. Journal of Nutrition 2003;133:2125-2130.

Vanharanta M, Voutilainen S, Rissanen T, et al: Risk of cardiovascular disease-related and all-cause death according to serum concentrations of enterolactone: Kuopio Ischaemic Heart Disease Risk Factor Study. Archives of Internal Medicine 2003;163:1099-1104.

Vioque J, Weinbrenner T, Castello A, et al: Intake of fruits and vegetables in relation to 10-year weight gain among Spanish adults. Obesity 2008;16:664-670.

Wang H, Nair M, Iezzoni G, et al: Quantification and characterization of anthocyanins in Balaton tart cherries. Journal of Agricultural and Food Chemistry 1997;45:2556-2560.

Wang H, Nair M, Strasburg G, et al: Antioxidant polyphenols from tart cherries (*Prunus cerasus*). Journal of Agricultural and Food Chemistry 1999;47:840-844.

Williams P, Grafenauer S, O'Shea J: Cereal grains, legumes, and weight management. Nutrition Reviews 2008;66:171-182.

Wu X, Beecher G, Holden J, et al: Lipophilic and hydrophilic antioxidant capacities of common foods in the United States. Journal of Agricultural and Food Chemistry 2004;52:4026-4037.

Youdim K, Dobbie M, Kuhnle G, et al: Interaction between flavonoids and the blood-brain barrier. Journal of Neurochemistry 2003;85:180-192.

Zemel M: Regulation of adiposity and obesity risk by dietary calcium: Mechanisms and implications. Journal of the American College of Nutrition 2002;21:146S-151S.

;b Chapter 6

Ahmad A, Greiner R, Moriguchi T, et al: Effects of dietary omega-3 fatty acid on the

morphology of the hippocampus. Journal of Neurochemistry 2000;74:88.

Arterburn L, Oken H, Hoffman J, et al: Algal-oil capsules and cooked salmon: nutritionally equivalent sources of docosahexaenoic acid. Journal of the American Dietetic Association 2008;108:1204-1209.

Arterburn L, Oken H, Hoffman J, et al: Bioequivalence of docosahexaenoic acid from different algal oils in capsules and in a DHA-fortified food. Lipids 2007;42:1011-1024.

Austria J, Richard M, Chahine M, et al: Bioavailability of alpha-linolenic acid in subjects after ingestion of three different forms of flaxseed. Journal of the American College of Nutrition 2008;27:214-221.

Bao D, Mori T, Burke V, et al: Effects of dietary fish and weight reduction on ambulatory blood pressure in overweight hypertensives. Hypertension 1998;32:710-717.

Bourre J: Dietary omega-3 fatty acids for women. Biomedical Pharmacotherapy 2007;61:105-112.

Bourre J: Dietary omega-3 fatty acids and psychiatry. Journal of Nutrition, Health, And Aging 2005;9:31-38.

Buydens-Branchev L, Branchev M, Hibbeln J: Associations between increases in plasma n-3 polyunsaturated fatty acids following supplementation and decreases in anger and anxiety in substance abusers. Progress in Neuropsychopharmacology 2008;32:568-575.

Chiu C, Huang S, Shen W, et al: Omega-3 fatty acids for depression in pregnancy. American Journal of Psychiatry 2003;160:385.

Conquer J, Holub B: Supplementation with an algae source of docosahexaenoic acid increases (n-3) fatty acid status and alters selected risk factors for heart disease in vegetarian subjects. Journal of Nutrition 1996;126:3032-3039.

Cott J, Hibbeln J: Lack of seasonal mood change in Icelanders. American Journal of Psychiatry 2001;158:328.

Delarue J, Matzinger O, Binnert C, et al: Fish oil prevents the adrenal activation elicited by mental stress in healthy men. Diabetes Metabolism 2003;29:289-295.

Esposito K, Pontillo A, di Palo C, et al: Effect of weight loss and lifestyle changes on vascular inflammatory markers in obese women. Journal of the American Medical Association 2003;289:1799-1804.

Freeman M, Hibbeln J, Wisner K, et al: Omega-3 fatty acids: Evidence basis for treatment and future research in psychiatry. Journal of Clinical Psychiatry 2006;67:1954-1967.

Haag M: Essential fatty acids and the brain. Canadian Journal of Psychiatry 2003;48:195-203.

Henriksen C, Haughholt K, Lindgren M, et al: Improved cognitive development among preterm infants attributable to early supplementation of human milk with docosahexaenoic acid and arachidonic acid. Pediatrics 2008;121:1137-1145.

Hibbeln J: From homicide to happiness: A commentary on omega-3 fatty acids in human Society. Nutrition & Health 2007;19:9-19.

Hibbeln J: Fish consumption and major depression. Lancet 1998;351:1213.

Hibbeln J, Nieminen L, Blasbalg T, et al: Healthy intakes of n-3 and n-6 fatty acids. American Journal of Clinical Nutrition 2006;83:1483S-1493S.

Horrocks L, Farooqui A: Docosahexaenoic acid in the diet: Its importance in maintenance and restoration of neural membrane function. Prostaglandins, Leukotrienes and Essential Fatty Acids 2004;70:361-372.

Huang T, Zandi P, Tucker K, et al: Benefits of fatty fish on dementia risk are stronger for those with APOE epsilon4. Neurology 2005;65:1409-1414.

Huang X, Xin X, McLennan P, et al: Role of fat amount and type in ameliorating diet-induced obesity. Diabetes, Obesity & Metabolism 2004;6:35-44.

International Society for the Study of Fatty Acids and Lipids: ISSFAL Newsletter 2007;Winter 14(3):1-13.

Kaur P, Hegglund I, Aschner M, et al: Docosahexaenoic acid may act as a neuroprotector for methylmercury-induced neurotoxicity in primary neural cell cultures. Neurotoxicology 2008;June 20th.

Kris-Etherton P, Hill A: n-3 fatty acids: Foods or supplements? Journal of the American Dietetic Association 2008;108:1125-1130.

Lim G, Calon F, Morihara T, et al: A diet enriched with the omega-3 fatty acid docosahexaenoic acid reduces amyloid burden in an aged Alzheimer mouse model. Journal of Neuroscience 2005;25:3042-3040.

Makrides M, Crowther C, Gibson R, et al: Docosahexaenoic acid and post-partum depression. Asia Pacific Journal of Clinical Nutrition 2003;12:S37.

Mamalakis G, Kalageropoulos N, Andrikopoulos N, et al: Depression and long chain n-3 fatty acids in adipose tissue in adults from Crete. European Journal of Clinical Nutrition 2006;60:882-888.

Mamalakis G, Tornaritis M, Kafatos A: Depression and adipose essential polyunsaturated fatty acids. Prostaglandins, Leukotrienes and Essential Fatty Acids 2002;67:311-318.

Molteni R, Barnard R, Ying Z, et al: A high-fat, refined sugar diet reduces hippocampal brain-derived neurotrophic factor, neuronal plasticity, and learning. Neuroscience 2002;112:803-814.

Mori T, Bao D, Burke V, et al: Dietary fish as a major component of a weight loss diet. American Journal of Clinical Nutrition 1999;70:817-825.

Morris M, Evans D, Bienias J, et al: Consumption of fish and n-3 fatty acids and risk of incident Alzheimer disease. Archives of Neurology 2003;60:940-946.

Motluk A: A fatty diet can clog your brain as well as your coronary arteries. New Science 2001;169:10.

Nitta H, Kinoyama m, Watanabe A, et al: Effects of nutritional supplementation with antioxidant vitamins and minerals and fish oil on antioxidant status and psychosocial stress in smokers. Clinical Experimental Medicine 2007;7:179-183.

Orr S, Bazinet R: The emerging role of docosahexaenoic acid in neuroinflammation. Current Opinions in Investigational Drugs 2008;9:735-743.

Owen C, Rees A, Parker G: The role of fatty acids in the development and treatment of mood disorders. Current Opinion in Psychiatry 2008;21:19-24.

Parker G, Gibson N, Brotchie H, et al: Omega-3 fatty acids and mood disorders. American Journal of Psychiatry 2006;163:969-978.

Parra D, Ramel A, Bandarra N, et al: A diet rich in long chain omega-3 fatty acids modulates satiety in overweight and obese volunteers during weight loss. Appetite 2008;June 14th.

Pjamondon H, Roberge M: Dietary PUFA supplements reduce memory deficits but not CA1 ischemic injury in rats. Physiology & Behavior 2008;95:492-500.

Pounis G, Panagiotakos D, Chrysohoou C, et al: Long-term fish consumption is associated with lower risk of 30-day cardiovascular disease events in survivors from an acute coronary syndrome. International Journal of Cardiology 2008;July 16th.

Roseland J, Haugen F, Hollung K, et al: Reduction of leptin gene expression by dietary polyunsaturated fatty acids. Journal of Lipid Research 2001;42:743-750.

Ruzickova J, Rossmeisl M, Prazak T, et al: Omega-3 PUFA of marine origin limit diet-induced

obesity in mice by reducing cellularity of adipose tissues. Lipids 2004;39:1177-1185.

Sanchez-Villegas A, Henriquez P, Figueiras A, et al: Long chain omega-3 fatty acids intake, fish consumption and mental disorders in the SUN cohort study. European Journal of Nutrition 2007;46:337-346.

Schaefer E, Bongard V, Beiser A, et al: Plasma phosphatidylcholine docosahexaenoic acid content and risk of dementia and Alzheimer disease. The Framingham Heart Study. Archives of Neurology 2006;63:1545-1550.

Schwellenbach L, Olson K, McConnell K, et al: The triglyceride-lowering effects of a modest dose of docosahexaenoic acid alone versus in combination with low dose eicosapentaenoic acid in patients with coronary artery disease and elevated triglycerides. Journal of the American College of Nutrition 2006;25:480-485.

Sinclair A, Begg D, Mathai M, et al: Omega 3 fatty acids and the brain. Asia Pacific Journal of Clinical Nutrition 2007;16 (suppl):391-397.

Somer E: The Origin Diet. New York, Owl Books, 2001, pp 48.

Sontrop J, Campbell M: Omega-3 polyunsaturated fatty acids and depression. Preventive Medicine 2006;42:4-13.

Stark K, Holub B: Differential eicosapentaenoic acid elevations and altered cardiovascular disease risk factor responses after supplementation with docosahexaenoic acid in postmenopausal women receiving and no receiving hormone replacement therapy. American Journal of Clinical Nutrition 2004;79:765-773.

Stevens L, Zentall S, Abate M, et al: Omega-3 fatty acids in boys with behavior, learning, and health problems. Physiology & Behavior 1996;59:915-920.

Stoll A, Locke C, Marangell L, et al: Omega-3 fatty acids and bipolar disorders. Prostaglandins,

Leukotrienes, and Essential Fatty Acids 1999;60:329-337.

Sublette M, Hibbeln J, Galfalvy H, et al: Omega-3 polyunsaturated essential fatty acid status as a predictor of future suicide risk. American Journal of Psychiatry 2006;163:1100-1102.

Tanaka K, Kon N, Ohkawa N, et al: Does breastfeeding in the neonatal period influence the cognitive function of very-low-birth-weight infants at 5 years of age? Brain & Development 2008; July 18th.

Tanskanen A, Hibbeln J, Tuomilehto J, et al: Fish consumption and depressive symptoms in the general population of Finland. Psychiatric Services 2001;52:529-531.

Timonen M, Horrobin D, Jokelainen J, et al: Fish consumption and depression. Journal of Affective Disorders 2004;82:447-452.

Tukiainen T, Tynkkynen T, Makinen V, et al: A multi-metabolite analysis of serum by (1)H NMR spectroscopy: Early systemic signs of Alzheimer's disease. Biochemical & Biophysical Research Communications 2008;375:356-361.

Virtanen J, Siscovick D, Longstreth W, et al: Fish consumption and risk of subclinical brain abnormalities on MRI in older adults. Neurology 2008;71:439-446.

Weldon S, Mullen A, Loscher C, et al: Docosahexaenoic acid induces an anti-inflammatory profile in lipopolysaccharide-stimulated human THP-1 macrophages more effectively than eicosapentaenoic acid. Journal of Nutritional Biochemistry 2007;18:250-258.

Whalley L, Fox H, Wahle K, et al: Cognitive aging, childhood intelligence, and the use of food supplements: Possible involvement of n-3 fatty acids. American Journal of Clinical Nutrition 2004;80:1650-1657.

;b Chapter 7

Baldewicz T, Goodkin K, Feaster D, et al: Plasma pyridoxine deficiency is related to increased psychological distress in recently bereaved homosexual men. Psychosomatic Medicine 1998;60:297-308.

Barragan L, Rodriguez M, Guerrero F: Depressive symptoms and hypomagnesemia in older diabetic subjects. Archives of Medical Research 2007;38:752-756.

Beard J, Hendricks M, Perez E, et al: Maternal iron deficiency anemia affects postpartum emotions and cognition. Journal of Nutrition 2005;135:267-272.

Bell I, Edman J, Morrow F, et al: Vitamin B1, B2, and B6 augmentation of tricyclic antidepressant treatment in geriatric depression with cognitive dysfunction. Journal of the American College of Nutrition 1992;11:159-163.

Bell I, Edman J, Morrow F, et al: B complex vitamin patterns in geriatric and young adult inpatients with major depression. Journal of the American Geriatric Society 1991;39:252-257.

Benton D, Cook R: The impact of selenium on mood. Biological Psychiatry 1991;29:1092-1098.

Benton D, Fordy J, Haller J: The impact of long-term vitamin supplementation on cognitive function. Psychopharmacology 1995;117:298-305.

Benton D, Haller J, Fordy J: Vitamin supplementation for 1 year improves mood. Neuropsychobiology 1995;32:98-105.

Bertone-Johnson E, Hankinson S, Bendich A, et al: Calcium and vitamin D intake and risk of incident premenstrual syndrome. Archives of Internal Medicine 2005;165:1246-1252.

Block G, Abrams B: Vitamin and mineral status of women of childbearing potential. Annals of the New York Academy of Sciences 1993;678:244-254.

Block G, Dresser C, Hartman A, et al: Nutrient sources in the American diet: Quantitative data from the NHANES. American Journal of Epidemiology 1985;122:13-26.

Bourre J: Effects of nutrients (in food) on the structure and function of the nervous system.

Journal of Nutrition and Health in Aging 2006;10:377-385.

Brocardo P, Budni J, Kaster M, et al: Folic acid administration produces an antidepressant-like effect in mice. Neuropharmacology 2008;54:464-473.

Brody S: High-dose ascorbic acid increases intercourse frequency and improves mood. Biological Psychiatry 2002;15:371-374.

Carroll D, Ring C, Suter M, et al: The effects of an oral multivitamin combination with calcium, magnesium, and zinc on psychological well-being in healthy young male volunteers.

Psychopharmacology 2000;150:220-225.

Chapuy M, Preziosi P, Maamer M, et al: Prevalence of vitamin D insufficiency in an adult normal population. Psychopharmacology 1998;135:319-323.

Clarke R, Birks J, Nexo E, Ureland P, et al: Low vitamin B12 status and risk of cognitive decline in older adults. American Journal of Clinical Nutrition 2007;86:1384-1391.

Dollahite J, Franklin D, McNew R: Problems encountered in meeting the Recommended Dietary Allowances for menus designed according to the Dietary Guidelines for Americans. Journal of the American Dietetic Association 1995;95:341-344.

Durga J, vanBoxtel M, Schouten E, et al: Effect of 3-year folic acid supplementation on cognitive function in older adults in the FACIT trial. Lancet 2007;369:208-216.

Eastley R, Wilcock G, Bucks R: Vitamin B12 deficiency in dementia and cognitive impairment. International Journal of Geriatric Psychiatry 2000;15:226-233.

Fatouti M, Paparrigopoulos T, Liappas J, et al: Mood disorder with mixed features due to vitamin B12 and folate deficiency. General Hospital Psychiatry 2002;24:106-109.

Gariballa S, Forster S: Effects of dietary supplements on depressive symptoms in older patients.

Clinical Nutrition 2007;26:545-551.

Gilbody S, Lightfoot T, Sheldon T: Is low folate a risk factor for depression? Journal of Epidemiology and Community Health 2007;61:631-637.

Gloth F, Alam W, Hollis B: Vitamin D vs broad spectrum phototherapy in the treatment of seasonal affective disorder. Journal of Nutrition and Health in Aging 1999;3:5-7.

Hampel J, Taylor C, Johnston C: Vitamin C deficiency and depletion in the United States: The Third National Health and Nutrition Examination Survey, 1988 to 1994. American Journal of Public Health 2004;94:870-875.

Harms L, Eyles D, McGrath J, et al: Developmental vitamin D deficiency alters adult behavior in mice. Behavioral Brain Research 2008;187:343-350.

Hathcock J, Shao A, Vieth R, et al: Risk assessment for vitamin D. American Journal of Clinical Nutrition 2007;85:6-18.

Hawkes W: The effect of selenium on mood in a confined human nutrition study. FASEB Journal 1994;8:A436.

Hvas A, Juui S, Bech P, et al: Vitamin B6 level is associated with symptoms of depression. Psychotherapy and Psychosomatics 2004;73:340-343.

Johnston C, Dancho C, Strong G: Orange juice ingestion and supplemental vitamin C are equally effective at reducing plasma lipid peroxidation in healthy adult women. Journal of the American College of Nutrition 2003;22:519-523.

Jorde R, Waterloo K, Saleh F, et al: Neuropsychological function in relation to serum parathyroid hormone and serum 25-hydroxyvitamin D levels. Journal of Neurology 2006;253:464-470.

Kaplan B, Crawford S, Gardner B, et al: Treatment of mood lability and explosive rage with minerals and vitamins. Journal of Child and Adolescent Psychopharmacology 2002;12:205-219.

Landsdowne A, Provost S: Vitamin D3 enhances mood in healthy subjects during winter.

Psychopharmacology 1998;135:319-323.

Lichtenstein A, Russell R: Essential nutrients: Food or supplements? Journal of the American

Medical Association 2005;294:351-358.

Liu B, Gordon M, Labranche J, et al: Seasonal prevalence of vitamin D deficiency in institutionalized older adults. Journal of the American Geriatric Society 1997;45:598-603.

Madabushi R, Frank B, Drewelow B, et al: Hyperforin in St. John's wort drug interactions.

European Journal of Clinical Pharmacology 2006;62:225-233.

Maes M, deVos N, Pioli R, et al: Lower serum vitamin E concentrations in major depression.

Journal of Affective Disorders 2000;58:241-246.

Matsuoka L, Wortsman J, Dannenberg M, et al: Clothing prevents ultraviolet-B radiation-dependent photosynthesis of vitamin D3. Journal of Clinical Endocrinology 1992;75:1099-1103.

Minasyan A, Keisala T, Lou Y, et al: Neurophobia, sensory and cognitive functions, and hedonic responses in vitamin D receptor mutant mice. Journal of Steroid Biochemistry and Molecular Biology 2007;104:274-280.

Mischoulon D: Update and critique of natural remedies as antidepressant treatments. Psychiatric Clinics of North America 2007;30:51-68.

Mischoulon D, Fava M: Role of S-adenosyl-methionine in the treatment of depression. American Journal of Clinical Nutrition 2002;76:1158S-1161S.

Nguyen M, Gregan A: S-adenosylmethionine and depression. Australian Family Physician 2002;31:339-342.

Owen A, Batterham M, Probst Y, et al: Low plasma vitamin E levels in major depression.

European Journal of Clinical Nutrition 2006;59:304-306.

Patterson A, Brown W, Roberts D: Dietary and supplement treatment of iron deficiency results in improvements in general health and fatigue in Australian women of childbearing age. Journal of the American College of Nutrition 2001;20:337-342.

Penninx B, Guralnik J, Ferrucci L, et al: Vitamin B12 deficiency and depression in physically disabled older women. American Journal of Psychiatry 2000;157:715-721.

Quaranta S, Buscaglia M, Meroni M, et al: Pilot study of the efficacy and safety of a modified-release magnesium 250 mg tablet (Sincromag) for the treatment of premenstrual syndrome. Clinical Drug Investigations 2007;27:51-58.

Rhainds M, Galibois I, Gagne D, et al: The effect of sunscreens use on vitamin D levels in young children. Epidemiology 2001;12:259.

Rugulies R: Depression as a predictor for coronary heart disease. American Journal of Preventive Medicine 2002;23:51-61.

Sarandol A, Sarandol E, Eker S, et al: Major depressive disorder is accompanied with oxidative stress. Human Psychopharmacology 2007;22:67-73.

Schneider B, Weber B, Frensch A, et al: Vitamin D in schizophrenia, major depression, and alcoholism. Journal of Neural Transmission 200;107:839-842.

Seelig M: Consequences of magnesium deficiency on the enhancement of stress reactions. Journal of the American College of Nutrition 1994;13:429-446.

Shea T: Effects of dietary supplementation with n-acetyl cysteine, acetyl-L carnitine and s-adenosyl methionine on cognitive performance and aggression in normal mice and mice expressing human ApoE4. Neuromolecular Medicine 2007;9:264-269.

Shikany J, Heimburer D, Piyathilake C, et al: Effect of folic acid fortification of foods on folate intake in female smokers with cervical dysplasia. Nutrition 2004;20:409-414.

Tavazzi B, Signoretti S, Lazzarino G, et al: Cerebral oxidative stress and depression of energy metabolism correlate with severity of diffuse brain injury in rats. Neurosurgery 2005;56:582-589.

The relationship between depression and serum ferritin level. European Journal of Clinical Nutrition 2007;61:532-535.

Thys Jacobs S, Starkey P, Bernstein D, et al: Calcium carbonate and the premenstrual syndrome: Effects on premenstrual and menstrual symptoms. American Journal of Obstetrics & Gynecology 1998;179:444-452.

Tolmunen T, Voutilainen S, Hintikka J, et al: Dietary folate and depressive symptoms are associated in middle-age Finnish men. Journal of Nutrition 2003;133:3233-3236.

Verdon F, Burnand B, Stubi C, et al: Iron supplementation for unexplained fatigue in non-anaemic women. British Journal of Medicine 2003;326:1124.

Vieth R: Why the optimal requirement for vitamin D3 is probably much higher than what is officially recommended for adults. Journal of Steroid Biochemistry and Molecular Biology 2004;89-90:575-579.

Vieth R: Vitamin D supplementation, 25-hydroxyvitamin D concentrations, and safety. American Journal of Clinical Nutrition 1999;69:842-856.

Vogiatzoglou A, Refsum H, Johnston C, et al: Vitamin B12 status and rate of brain volume loss in community-dwelling elderly. Neurology 2008;71:826-832.

Walker A, deSouza M, Vickers M, et al: Magnesium supplementation alleviates premenstrual syndromes of fluid retention. Journal of Womens Health 1998;7:1157-1165.

Widmer J, Henrotte J, Raffin Y, et al: Relationship between erythrocyte magnesium, plasma electrolytes and cortisol, and intensity of symptoms in major depressed patients. Journal of Affective Disorders 1995;34:201-209.

Wilkins C, Sheline Y, Roe C, et al: vitamin D deficiency is associated with low mood and worse cognitive performance in older adults. American Journal of Geriatric Psychiatry 2006;14:1032-1040.

Williams A, Cotter A, Sabina A, et al: The role of vitamin B6 as treatment for depression: A systematic review. Family Practice 2005;22:532-537.

Wouters W, Wagenaar L, Rozendaal M, et al: Effect of an enriched drink on cognitive function in frail elderly persons. Journals of Gerontology 2005;60:265-270.

Wyatt K, Dimmock P, Jones P, et al: Efficacy of vitamin B6 in the treatment of premenstrual syndrome: Systematic review. British Medical Journal 1999;318:1375-1381.

Zandi P, Anthony J, Khachaturian A, et al: Reduced risk of Alzheimer disease in users of antioxidant vitamin supplements. Archives of Neurology 2004;61:82-88.

Ziambaras K, Jack S: Reversible muscle weakness in patients with vitamin D deficiency. Western Journal of Medicine 1997;167:435-439.

Zittermann A: Vitamin D in preventive medicine: Are we ignoring the evidence? British Journal of Nutrition 2003;89:552-572.

;b Chapter 8

Acheson K: Caffeine and insulin sensitivity. Metabolic Syndrome and Related Disorders 2005;3:19-25.

Berger R, Shaw J, Decatanzaro D: Impact of acute bisphenol-A exposure upon intrauterine implantation of fertilized ova and urinary levels of progesterone and 17beta-estradiol.

Reproductive Toxicology 2008;26:94-99.

Buemann B, Toubro S, Astrup A: The effect of wine or beer versus a carbonated soft drink,

served at a meal. On ad libitum energy intake. International Journal of Obesity 2002;26:1367-1372.

Chevrier J, Eskenazi B, Holland N, et al: Effects of exposure to polychlorinated biphenyls and organochlorine pesticides on thyroid function during pregnancy. American Journal of Epidemiology 2008;168:298-310.

Christensen L, Burrows R: Dietary treatment of depression. Behavioral Therapy 1990;21:183-193.

Christensen L, Krietsch K, White B, et al: Impact of a dietary change on emotional distress. Journal of Abnormal Psychiatry 1985;94(4):565-579.

Christensen L, Pettijohn L: Mood and carbohydrate cravings. Appetite 2001;36:137-145.

DiMeglio D, Mattes R: Liquid versus solid carbohydrate: Effects on food intake and body weight. International Journal of Obesity 2000;24:794-800.

Dubois L, Farmer A, Girard M, et al: Regular sugar-sweetened beverage consumption between meals increases risk of overweight among preschool-aged children. Journal of the American Dietetic Association 2007;107:924-934.

Guéguen N, Jacob C, Le Guellec H, et al: Sound Level of Environmental Music and Drinking Behavior: A Field Experiment With Beer Drinkers. Alcoholism: Clinical and Experimental Research 2008;32:1795-1798.

Harrington S: The role of sugar-sweetened beverage consumption in adolescent obesity. Journal of School Nursing 2008;24:3-12.

Hetherington M, Cameron F, Wallis D, et al: Stimulation of appetite by alcohol. Physiology & Behavior 2001;74:283-289.

Holt S, Sandona N, Brand-Miller J: The effects of sugar-free vs sugar-rich beverages on feelings

of fullness and subsequent food intake. International Journal of Food and Sports Nutrition 2000;51:59-71.

Kerr W, Patterson D, Koenen M, et al: Alcohol Content Variation of Bar and Restaurant Drinks in Northern California. Alcoholism: Critical and Experimental Research 2008;32:1623-1629.

Lands W: Alcohol, calories, and appetite. Vitamins and Hormones 1998;54:31-49.

Lang I, Galloway T, Scarlett A, et al: Association of urinary bisphenol A concentration with medical disorders and laboratory abnormalities in adults. Journal of the American Medical Association 2008;300:1303-1310.

Leranth C, Hajszan T, Szigeti-Buck J, et al: Bisphenol A prevents the synaptogenic response to estradiol in hippocampus and prefrontal cortex of ovariectomized nonhuman primates. Proceedings of the National Academy of Sciences 2008;105:14187-14191.

Lieberman H: Hydration and cognition. Journal of the American College of Nutrition 2007;555S-561S.

Ludwig D, Peterson K, Gortmaker S: Relation between consumption of sugar-sweetened drinks and childhood obesity. Lancet 2001;357:505-508.

Malik V, Schulze M, Hu F: Intake of sugar-sweetened beverages and weight gain. American Journal of Clinical Nutrition 2006;84:274-288.

Mattes R, Campbell W: Effects of food form and timing of ingestion on appetite and energy intake in lean young adults and in young adults with obesity. Journal of the American Dietetic Association 2009;109:430-437.

Mattes R, Rothacker D: Beverage viscosity is inversely related to postprandial hunger in humans. Physiology & Behavior 2001;74:551-557.

Odegaard A, Pereira M, Koh W, et al: Coffee, tea, and incident type 2 diabetes. American Journal

of Clinical Nutrition 2008;88:979-985.

Lopez-Garzia E, van Dam R, Li T, et al: The relationship of coffee consumption with mortality.

Annals of Internal Medicine 2008;148:904-914.

Raloff J: Coffee, spices, and wine. Science News 2004;165:282-284.

Rolls B, Bell E, Thorwart M: Water incorporated into a food but not served with a food decreases energy intake in lean women. American Journal of Clinical Nutrition 1999;70:448-455.

Schulze M, Manson J, Ludwig D, et al: Sugar-sweetened beverages, weight gain, and incidence of type 2 diabetes in young and middle-aged women. Journal of the American Medical Association 2004;292:927-934.

;b Chapter 9

Barger J, Kayo T, Vann J, et al: A low dose of dietary resveratrol partially mimics caloric restriction and retards aging parameters in mice. PloS ONE 2008;308:e2264.

Buijsse B, Feskens E, Kok F, et al: Cocoa intake, blood pressure, and cardiovascular mortality.

The Zutphen Elderly Study. Archives of Internal Medicine 2006;166:411-417.

Cao G, Sofie E, Prior R: Antioxidant capacity of tea and common vegetables. Journal of Agricultural Food Chemistry 1996;44:3426-3431.

Dillinger T, Barriga P, Escarcega S, et al: Food of the Gods: Cure for humanity? Journal of Nutrition 2000;130:2057-2072.

Ding E, Hutfless S, Ding X, et al: Chocolate and prevention of cardiovascular disease. Nutrition & Metabolism 2006;3:2.

Dulloo A, Duret C, Rohrer D, et al: Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans. American Journal

of Clinical Nutrition 1999;70:1040-1045.

Dulloo A, Seydoux J, Girardier L, et al: Green tea and thermogenesis: interactions between catechin-polyphenols, caffeine, and sympathetic activity. International Journal of Obesity 2000;24:252-258.

Fletcher B, Pine K, Woodbridge Z, et al: how visual images of chocolate affect the craving and guilt of female dieters. Appetite 2007;48:211-217.

Giu L, House S, Wu X, et al: Procyanidin and catechin contents and antioxidant capacity of cocoa and chocolate products. Journal of Agricultural Food Chemistry 2006;54:4057-4061.

Halsted C, Villanueva J, Devlin A, et al: Metabolic interactions of alcohol and folate. Journal of Nutrition 2002;132:2367-2372.

Heinrich U, Neukam K, Tronnier H, et al: Long-term ingestion of high flavonol cocoa provides photoprotection against UV-induced erythema and improves skin condition in women. Journal of Nutrition 2006;136:1565-1569.

Hetherington M, Boyland E: Short-term effects of chewing gum on snack intake and appetite. Appetite 2007;48:397-401.

Hetherington M, MacDiarmid J: Pleasure and excess: Liking for and overconsumption of chocolate. Physiology & Behavior 1995;57:27-35.

Kao Y, Hiipakka R, Liao S: Modulation of obesity by a green tea catechin. American Journal of Clinical Nutrition 2000;72:1232-1233.

Karim M, McCormick K, Kappagoda T: Effects of cocoa extracts on endothelium-dependent relaxation. Journal of Nutrition 2000;130:2105-2108.

Keen C, Holt R, Oteiza P, et al: Cocoa antioxidants and cardiovascular health. American Journal of Clinical Nutrition 2005;81:298S-303S.

Kris-Etherton P, Derr J, Mustad V, et al: Effects of a milk chocolate bar per day substituted for a high-carbohydrate snack in young men on an NCEP/AHA Step 1 diet. American Journal of Clinical Nutrition 1994;60:1037S-1042S.

Lee K, Kim Y, Lee H, et al: Cocoa has more phenolic phytochemicals and a higher antioxidant capacity than teas and red wine. Journal of Agricultural Food Chemistry 2003;51:7292-7295.

McKay D, Blumberg J: The role of tea in human health: An update. Journal of the American College of Nutrition 2002;21:1-13.

Mehlig K, Skoog I, Guo X, et al: Alcoholic beverages and incidence of dementia. American Journal of Epidemiology 2008;167:684-691.

Miller K, Stuart D, Smith N, et al: Antioxidant activity and polyphenol and procyanidin contents of selected commercially available cocoa-containing chocolate products in the United States. Journal of Agricultural Food Chemistry 2006;54:4062-4068.

Ng T, Feng L, Niti M, et al: Tea consumption and cognitive impairment and decline in older Chinese adults. American Journal of Clinical Nutrition 2008;88:224-231.

Rein D, Paglieroni T, Pearson D, et al: Cocoa and wine polyphenols modulate platelet activation and function. Journal of Nutrition 2000;130:2120-2126.

Rossner S: Chocolate: divine food, fattening junk, or nutritious supplementation? European Journal of Clinical Nutrition 1997;51:341-345.

Rumpler W, Seale J, Clevidence B, et al: Oolong tea increases metabolic rate and fat oxidation in men. Journal of Nutrition 2001;131:2848-2852.

Soleas G, Diamandis E, Goldberg D: Wine as a biological fluid: History, production, and role in disease prevention. Journal of Clinical Lab An 1997;11:287-313.

Stephoe A, Gibson E, Vuononvirta R, et al: The effects of tea on psychophysiological stress

responsivity and post-stress recovery. Psychopharmacology 2007;190:81-89.

Wan Y, Vinson J, Etherton T, et al: Effects of cocoa powder and dark chocolate on LDL oxidative susceptibility and prostaglandin concentrations in humans. American Journal of Clinical Nutrition 2001;74:596-602.

Wiseman S, Balentine D, Frei B; Antioxidants in tea. Critical Reviews in Food Science and Nutrition 1997;37:705-718.

Wu, Lu, Chang C, et al: Relationship among habitual tea consumption, percent body fat, and body fat distribution. Obesity Research 2003;11:1088-1095.

;b Chapter 10

Akata T, Sekiguchi S, Takahashi M, et al: Successful combined treatment with vitamin B12 and bright artificial light of one case with delayed sleep phase syndrome. Japanese Journal of Psychiatry and Neurology 1993;47:439-440.

Baumann P, Steinberg R, Koeb I, et al: Long-term treatment of hyposomnia with L-tryptophan. Experimentia 1985;41:1214.

Brown S, Salive M, Pahor M, et al: Occult caffeine as a source of sleep problems in an older population. Journal of the American Geriatric Society 1995;43:860-864.

Cheruku S, Montgomery-Downs H, Farkas S, et al: Higher maternal plasma docosahexaenoic acid during pregnancy is associated with more mature neonatal sleep-state patterning. American Journal of Clinical Nutrition 2002;76:608-613.

Chollet D, Franken P, Raffin Y, et al: Magnesium involvement in sleep. Behavior Genetics 2001;31:413-425.

Dimpfel W, Suter A: Sleep improving effects of a single dose administration of a valerian/hops fluid extract. European Journal of Medical Research 2008;13:200-204.

Dixon J, Schachter L, O'Brien P: Sleep disturbance and obesity. Archives of Internal Medicine 2001;16:102-106.

Donath F, Quispe S, Diefenbach K, et al: Critical evaluation of the effect of valerian extract on sleep structure and sleep quality. Pharmacopsychiatry 2000;33:47-53.

Fitzgerald M: Restless leg syndrome: Iron important. The Nurse Practitioner 2008;33:7-9.

Guan Z, Vgontzas A, Bixler E, et al: Sleep is increased by weight gain and decreased by weight loss in mice. Sleep 2008;31:627-633.

Hajak G, Huether G, Blanke J, et al: The influence of intravenous L-tryptophan on plasma melatonin and sleep in men. Pharmacopsychiatry 1991;24:17-20.

Lavialle M, Champeil-Potokar G, Alessandri J, et al: An (n-3) polyunsaturated fatty acid-deficient diet disturbs daily locomotor activity, melatonin rhythm, and striatal dopamine in Syrian hamsters. Journal of Nutrition 2008;138:1719-1724.

Leathwood P, Chauffard F, Heck E, et al: Aqueous extract of valerian root (*Valeriana officinalis* L.) improves sleep quality in man. Pharmacology, Biochemistry and Behavior 1982;17: 65-71.

Llanas A, Hachul H, Bittencourt L, et al: Physical therapy reduces insomnia symptoms in postmenopausal women. Maturitas 2008;September 24th.

Ohta T, Ando K, Iwata T, et al: Treatment of persistent sleep-wake schedule disorders in adolescents with methylcobalamine (vitamin B12). Sleep 1991;14:414-418.

Okawa M, Mishima K, Nanami T, et al: Vitamin B12 treatment for sleep wake rhythm disorders. Sleep 1990;13:15-23.

Pollet P, Leathwood P: The influence of tryptophan on sleep in man. International Journal of Vitamin Nutrition Research 1983;53:223.

Spinweber C: L-tryptophan administered to chronic sleep-onset insomniacs.

Psychopharmacology 1986;90:151-155.

Su K: Mind-body interface: The role of n-3 fatty acids in psychoneuroimmunology, somatic presentation, and medical illness comorbidity of depression. Asia Pacific Journal of Clinical Nutrition 2008;17 (suppl):151-157.

Waller S, Vander J, Klurfeld D, et al: Evening ready-to-eat cereal consumption contributes to weight management. Journal of the American College of Nutrition 2004;23:316-321.

Yehuda S, Rabinovitz S, Mostofsky D: Essential fatty acids and sleep: Mini-review and hypothesis. Medical Hypothesis 1998;50:139-145.

;b Chapter 11

Augestad L, Slettemoen R, Flanders W: Physical activity and depressive symptoms among Norwegian adults aged 20-50. Public Health Nursing 2008;25:536-545.

Avila-Funes J, Gray-Donald K, Payette H: Association of nutritional risk and depressive symptoms with physical performance in the elderly. Journal of the American College of Nutrition 2008;27:492-498.

Blumenthal J, Babyak M, Doraiswamy P, et al: Exercise and pharmacotherapy in the treatment of major depressive disorder. Psychosomatic Medicine 2007;69:587-596.

Blumenthal J, Babyak M, Moore K, et al: Effects of exercise training on older patients with major depression. Archives of Internal Medicine 1999;159:2349-2356.

Bots S, Tijhuis M, Giampaoli S, et al: Lifestyle- and diet-related factors in late-life depression. International Journal of Geriatric Psychiatry 2007;23:478-484.

Cherkas L, Hunkin J, Kato B, et al: The association between physical activity in leisure time and

leukocyte telomere length. Archives of Internal Medicine 2008;168:154-158.

Choi P, Van Horn J, Picker D, et al: Mood changes in women after an aerobics class. Health Care for Women International 1993;14:167-177.

Conroy M, Simkin-Silverman L, Pettee K, et al: Lapses and psychosocial factors related to physical activity in early postmenopause. Medicine and Sciences in Sports Exercise 2007;39:1858-1866.

Coyle C, Santiago M: Aerobic exercise training and depressive symptomatology in adults with physical disabilities. Archives of Physical Medicine and Rehabilitation 1995;76:647-652.

Desharnais R, Jobin J, Cote C, et al: Aerobic exercise and the placebo effect. Psychosomatic Medicine 1993;55:149-154.

Folkens C: Effects of physical training on mood. Journal of Clinical Psychology 1976;32:385-388.

Foreyt J, Brunner R, Goodrick G, et al: Psychological correlates of reported physical activity in normal-weight and obese adults. International Journal of Obesity 1995;19:S69-S72.

Goldfarb A, Jamurtas A: Beta-endorphin response to exercise. Sports Medicine 1997;24:8-16.

Harber V, Sutton J, MacDougall J, et al: Plasma concentrations of beta-endorphin in trained eumenorrheic and amenorrheic women. Fertility and Sterility 1997;67:648-653.

Heitkamp H, Huber W, Scheib K: Beta-endorphin and adrenocorticotropin after incremental exercise and marathon running. European Journal of Applied Physiology 1996;72:417-424.

King A, Haskell W, Taylor C, et al: Group- vs home-based exercise training in healthy older men and women. Journal of the American Medical Association 1991;266:1535-1542.

King A, Taylor C, Haskell W: Effects of differing intensities and formats of 12 months of exercise training on psychological outcomes in older adults. Health Psychology 1993;12:292-300.

Lautenschlager N, Cox K, Flicker L, et al: Effect of physical activity on cognitive function in

older adults at risk for Alzheimer disease. Journal of the American Medical Association 2008;300:1027-1037.

Matthews C, Chen K, Freedson P, et al: Amount of time spent in sedentary behaviors in the United States, 2003-2004. American Journal of Epidemiology 2008;167:875-881.

Merrill R, Aldana S, Greenlaw R, et al: The effects of an intensive lifestyle modification program on sleep and stress disorders. Journal of Nutrition, Health & Aging 2007;11:242-248.

Nelson D, Sammel M, Freeman E, et al: Effect of physical activity on menopausal symptoms among urban women. Medicine & Science in Sports & Exercise 2008;40:50-58.

Penaskovic K, Baker P, Bamman M, et al: Depressive symptoms and physical activity in older adults. Journal of the American Geriatric Society 2002;50:P233.

Phelan S, Roberts M, Lang W, et al: Empirical evaluation of physical activity recommendations for weight control in women. Medicine and Science in Sports Exercise 2007;39:1832-1836.

Pinchasov B, Shurgaja A, Grischin O, et al: Mood and energy regulation in seasonal and non-seasonal depression before and after midday treatment with physical exercise or bright light. Psychiatry Research 2000;94:29-42.

Ruuskanen J, Fuoppila I: Physical activity and psychological well-being among people aged 65 to 84 years. Age and Ageing 1995;24:292-296.

Schwarz L, Kindermann W: Changes in beta-endorphin levels in response to aerobic and anaerobic exercise. Sports Medicine 1992;13:25-36.

Sorensen M, Anderssen S, Hjerman I, et al: Exercise and diet interventions improve perceptions of self in middle-aged adults. Scandinavian Journal of Medicine & Science in Sports 1997;7:312-320.

Sorensen M, Anderssen S, Hjerman I, et al: The effect of exercise and diet on mental health and

quality of life in middle-aged individuals with elevated risk factors for cardiovascular disease.

Journal of Sports Science 1999;17:369-377.

Teychenne M, Ball K, Salmon J: Physical activity and likelihood of depression adults. Preventive

Medicine 2008;46:397-411.